# Recovery of African wild dogs suppresses prey but does not trigger a trophic cascade

Adam T. Ford,<sup>1,2,9</sup> Jacob R. Goheen,<sup>2,3</sup> David J. Augustine,<sup>4</sup> Margaret F. Kinnaird,<sup>2,5</sup> Timothy G. O'Brien,<sup>2,5</sup> Todd M. Palmer,<sup>2,6</sup> Robert M. Pringle,<sup>2,7</sup> and Rosie Woodroffe<sup>2,8</sup>

> <sup>1</sup>Department of Zoology, University of British Columbia, Vancouver, British Columbia V6T1Z4 Canada <sup>2</sup>Mpala Research Centre, Nanyuki, Kenya

<sup>3</sup>Department of Zoology and Physiology, University of Wyoming, Laramie, Wyoming 82071 USA

<sup>4</sup>Rangeland Resources Research Unit, USDA-ARS, Fort Collins, Colorado 80526 USA

<sup>5</sup>Wildlife Conservation Society, Global Conservation Programs, Bronx, New York 10460 USA

<sup>6</sup>Department of Biology, University of Florida, Gainesville, Florida 32611 USA

<sup>7</sup>Department of Ecology and Evolutionary Biology, Princeton University, Princeton, New Jersey 08544 USA <sup>8</sup>Institute of Zoology, Zoological Society of London Regent's Park, London NW14RY United Kingdom

*Abstract.* Increasingly, the restoration of large carnivores is proposed as a means through which to restore community structure and ecosystem function via trophic cascades. After a decades-long absence, African wild dogs (Lycaon pictus) recolonized the Laikipia Plateau in central Kenya, which we hypothesized would trigger a trophic cascade via suppression of their primary prey (dik-dik, Madoqua guentheri) and the subsequent relaxation of browsing pressure on trees. We tested the trophic-cascade hypothesis using (1) a 14-year time series of wild dog abundance; (2) surveys of dik-dik population densities conducted before and after wild dog recovery; and (3) two separate, replicated, herbivore-exclusion experiments initiated before and after wild dog recovery. The dik-dik population declined by 33% following wild dog recovery, which is best explained by wild dog predation. Dik-dik browsing suppressed tree abundance, but the strength of suppression did not differ between before and after wild dog recovery. Despite strong, top-down limitation between adjacent trophic levels (carnivoreherbivore and herbivore-plant), a trophic cascade did not occur, possibly because of a time lag in indirect effects, variation in rainfall, and foraging by herbivores other than dik-dik. Our ability to reject the trophic-cascade hypothesis required two important approaches: (1) temporally replicated herbivore exclusions, separately established before and after wild dog recovery; and (2) evaluating multiple drivers of variation in the abundance of dik-dik and trees. While the restoration of large carnivores is often a conservation priority, our results suggest that indirect effects are mediated by ecological context, and that trophic cascades are not a foregone conclusion of such recoveries.

Key words: Acacia; African wild dogs (Lycaon pictus); antelope; dik-dik (Madoqua guentheri); endangered species; food web; indirect effect; large carnivore; rain; savanna; tree cover.

#### INTRODUCTION

Carnivores can powerfully shape ecosystems through their direct effect on herbivores, and their resulting indirect effect on plants and abiotic processes such as nutrient cycling, erosion, and fire (e.g., Hairston et al. 1960, Estes et al. 1998, Schmitz et al. 2004, Croll et al. 2005, Estes et al. 2011). The strength of these indirect effects has been used to justify conservation efforts, with the prediction that the restoration of large carnivores will trigger a trophic cascade (Mech 2012, Ripple et al. 2014). Ecologists have struggled to quantify this prediction, and so there remains a number of unresolved questions both about the strength and generality of topdown control as well as the mechanisms by which large carnivores indirectly benefit plants (Kauffman et al. 2010, Kuker and Barrett-Lennard 2010, Estes et al. 2011, Beschta and Ripple 2012, 2013, Mech 2012, Winnie 2012, 2014, Newsome et al. 2013, Beschta et al. 2014, Peterson et al. 2014). Thus, while we know that large carnivores can affect important ecosystem processes in some cases, the question remains: in which ecological contexts do the indirect effects of carnivores exert primacy over other drivers of species abundance?

Terrestrial food webs are embedded within complex and shifting ecological contexts that determine the strength of indirect effects (Schmitz 2010). This context may include the presence of reticulate food chains, donor control, and environmental heterogeneity (Strong 1992, Polis and Strong 1996, Polis et al. 2000). Reticulate food chains encompass multiple species with similar resource requirements within a given trophic

Manuscript received 3 November 2014; revised 20 February 2015; accepted 25 March 2015; final version received 15 April 2015. Corresponding Editor: M. Festa-Bianchet.

<sup>&</sup>lt;sup>9</sup> Present address: Department of Integrative Biology, University of Guelph, Guelph, Ontario N1G 2W1 Canada. E-mail: adamford@uoguelph.ca

level (Polis and Strong 1996, Tschanz et al. 2007, Thibault et al. 2010). Following the decline of a single species of consumer, functional or numerical compensation within that trophic level may buffer against a trophic cascade (Finke and Denno 2004). For example, wolves, grizzly bears, and cougars may all contribute to the decline of elk and the release of aspen in the Greater Yellowstone Ecosystem, muddling causation from any single predator (Peterson et al. 2014). Donor control arises when organisms defend themselves (e.g., secondary compounds or defensive armaments) or otherwise impede (e.g., risk-avoidance behavior in animals) the flow of energy to higher trophic levels within food chains (Polis and Strong 1996, van der Stap et al. 2007, Mooney et al. 2010). For example, impala avoid risky areas of the landscape, leading to the suppression of preferred plants and the domination of well-defended plants in safe areas (Ford et al. 2014). Environmental heterogeneity, particularly variation in light, soil nutrients, and rainfall, can limit plant abundance more than herbivory (Leibold 1989, Schmitz 1994). Lack of rainfall can reduce resource availability for herbivores, thereby limiting populations directly (Hopcraft et al. 2010) or increasing the vulnerability of individuals to predation (Sinclair and Arcese 1995). Together, reticulate food chains, donor control, and environmental heterogeneity shape the ecological context in which trophic cascades either emerge or are overridden in terrestrial food webs.

We tested the trophic-cascade hypothesis in Laikipia, Kenya, a 12000-km<sup>2</sup> region that was naturally recolonized by African wild dogs (Lycaon pictus) following a 20-year absence (Woodroffe 2011). About 60% of African wild dog diets are composed of dik-dik (Madoqua guentheri; Woodroffe et al. 2007), which is also the most abundant ungulate in this region (Augustine 2010). Previous work in this system indicates that herbivory by small-sized (i.e., dik-dik) and mediumsized (i.e., impala) ungulates limit the biomass of tree communities (Augustine and McNaughton 2004, Goheen et al. 2013, Ford et al. 2014). Given the importance of dik-dik as prey for wild dogs and the potential effect of dik-dik on tree abundance, there is potential that wild dog recovery triggered a density-mediated trophic cascade. However, reticulate food chains, donor control, and environmental heterogeneity are also present in this system: both wild dogs and dik-dik coexist alongside a diverse assemblage of competitors; savanna ecosystems are characterized by unstable variation in rainfall that limits the distribution of tree cover (Sankaran et al. 2005); trees consumed by dik-dik possess chemical and mechanical defenses that can alter the direction of trophic cascades (Ford et al. 2014). Thus, in addition to a trophic cascade, we evaluated multiple sources of causality that may also explain variation in the abundance of dik-dik and trees.

Specifically, we assembled data to test the following predictions: (1) that wild dogs suppress the abundance of dik-dik; (2) that dik-dik are capable of suppressing

the abundance of trees; and (3) that the effect of dik-dik on tree abundance was reduced in the presence of wild dogs. To test these predictions, we monitored wild dog and dik-dik populations for 14 years, and used size selective ungulate-exclusion plots to quantify the effect of herbivory by dik-dik. A separate set of exclusion plots was established both before and after wild dog recovery, and therefore enabled us to test whether predation by wild dogs decreased the net effect of herbivory on tree abundance.

#### Methods

#### Prediction 1: Wild dogs suppress dik-dik abundance

Since their return to the study area in 2002, we have monitored wild dogs at Mpala Research Centre (MRC) using global positioning system (GPS) telemetry and radio-telemetry to quantify pack-level biomass. We monitored the abundance (i.e., density) of dik-dik using distance sampling methods on a semiannual basis from 1999 to 2002 and again from 2008 to 2014. Details on the study area and methods for monitoring wild dogs and dik-dik are provided in Appendix A.

We evaluated four lines of evidence to assess how wild dogs affected the dik-dik population. First, we compared the density (individuals/km<sup>2</sup>) of dik-dik before and after wild dog recovery using a generalized least squares (GLS) analysis. We used a GLS because of nonindependence between sequential estimates of density. For this and all subsequent GLS analyses, we tested for serial autocorrelation of residuals using autocorrelation and partial autocorrelation functions. Following Zuur et al. (2009), we incorporated both correlation and variance structures into the model and present coefficient estimates based on restricted maximum likelihood estimation. A summary of these models is provided in Appendix A.

Second, we quantified the effect of the estimated consumption of dik-dik by wild dogs on the population growth rate (r) of dik-dik. We estimated consumption of dik-dik based on the energetic demand of wild dogs combined with the energetic return of an adult dik-dik (Woodroffe et al. 2007). An average-sized wild dog (25.2 kg) that fed hypothetically and exclusively on dik-dik would require the caloric return of 0.61 dik-dik per day; however, because dik-dik account for  $\sim 62\%$  of prey biomass of wild dogs at our study site (Woodroffe et al. 2007), the predicted demand of dik-dik for an averagesized wild dog is 0.378 dik-dik per day, or 0.015 dik-dik per day per kilogram of wild dog. Thus, to estimate consumption of dik-dik by wild dogs, we multiplied 0.015 by the estimated biomass of wild dog packs on MRC and by the number of days each pack spent in the area. To quantify the population growth rate of dik-dik, we calculated

$$r=\frac{N_{i+1}-N_i}{t_{i+1}-t_i},$$

where N is the population density estimate of dik-dik

from the *i*th survey at time t. We then used r as the response variable in a GLS regression, and the estimated consumption of dik-dik by wild dogs between  $t_i$  and  $t_{i+1}$ as a predictor. We assumed the number of dik-dik consumed per kilogram of wild dog has remained constant among all dik-dik population surveys. However, foraging theory suggests that the rate of dik-dik consumption by wild dogs may change with the density of dik-dik (i.e., a Type I, II, or III functional response [Holling 1959]). If the functional response of wild dogs changed, then the estimated consumption of dik-dik by wild dogs would interact with dik-dik population density to affect r. We therefore tested an interaction between the estimated consumption of dik-dik by wild dogs and the population density of dik-dik on r. We also tested for density dependence in the dik-dik population using  $N_i$  as a predictor variable and r as the response. If the dik-dik population is experiencing density-dependent growth, then  $N_i$  will have a negative effect on r; such density dependence could confound the potential effects of wild dog recovery on the dik-dik population. We assessed the effects of wild dog predation, rainfall, and density dependence using an information-theoretic approach, and Akaike information criterion corrected for small sample sizes (AIC<sub>c</sub>) to evaluate support for competing models (see Appendix B).

Third, we quantified the effect of the estimated consumption of dik-dik by wild dogs on an index of dik-dik recruitment. Following Augustine (2010), we used the proportion of dik-dik groups consisting of three or more individuals for each population survey as an index of recruitment. Because dik-dik are territorial, monogamous, and females typically give birth to a single offspring, the presence of a group of three is almost always the result of successful reproduction (Kingswood and Kumamoto 1996, Komers 1996). We used a GLS regression to test for the effect of consumption by wild dogs on the recruitment index.

Fourth, a wild dog den was established midway through a series of line-transect surveys conducted in 2011-2012, and we quantified short-term responses of dik-dik to this event. The den was established in December 2011, occupied by 31 individuals (19 adults, 12 pups), and was abandoned after the pups were fully weaned in late January 2012. While denning, wild dogs typically increase their consumption of dik-dik by 10% (Woodroffe et al. 2007) and forage almost exclusively within 3 km of the den site (i.e., the denning home range; Appendix B: Fig. B1). This shift in the diet and movements associated with wild-dog denning allowed us to investigate responses of dik-dik to a short-term pulse of intense predation. Our 2011-2012 surveys were conducted in addition to the surveys used to estimate dik-dik density across MRC and were focused on a subsection of the study area. We quantified encounter rates (number of dik-dik/km) with dik-dik along a 14km road transect before (pre-denning, November 2011), during (active denning, January 2012), and after (postdenning, March 2012) this den was used by wild dogs. Because dik-dik are territorial, we do not expect that short-term changes in abundance would be caused by emigration of dik-dik from the denning home range. We compared dik-dik encounter rates in the denning home range to a 17-km transect in a nearby control area where wild dogs did not forage as frequently while the den was active (Appendix B: Fig. B1). The control area consisted of similar habitat and climate as the denning home range, and was accessible to wild dogs based on our telemetry study. To quantify encounter rates, we drove 10 km/h with two dedicated observers and one driver to locate dik-dik. Typically, distance sampling methods are preferred to encounter rates because the former provides an estimate of variance and accounts for non-detection. However, due to lack of temporal replication of transects within each pre-, active-, and post-denning survey period, density estimates could not be derived from these surveys. We validated the relationship between encounter rates and density in our study area using the 16 population surveys conducted across MRC (i.e., excluding the surveys conducted in the control and denning areas in 2011–2012) between 1999 and 2013 ( $r^2$ = 0.873, P < 0.0001, indicating that encounter rates provide an accurate index of dik-dik density. We used an exact test with a Poisson distribution to evaluate the null hypothesis that the encounter rate during the activedenning period did not differ from that of the predenning or post-denning periods. We performed separate exact tests for the denning home range and the control area. In addition to encounter rates, we also compared the recruitment index in the denning and control areas among pre-, active-, and post-denning survey periods using a proportion test. This test evaluates the null hypothesis that recruitment does not change with the denning activity of wild dogs. We performed separate proportion tests for the denning home range and the control area.

#### Other potential drivers of dik-dik abundance

We considered three possible alternatives to wild dog recovery that may explain variation in dik-dik density. First, we evaluated whether populations of other large carnivores had increased along with wild dogs, thereby contributing to the suppression of dik-dik and confounding the effect of wild dog recovery. We focused on species of carnivore likely to consume a significant number of dik-dik (i.e., leopards [Panthera pardus] and black-backed jackals [Canis mesomelas] [Estes 1991]) and compared the number of detections from a cameratrapping survey conducted before wild dog recovery (2000-2002; 7364 trap hours at 19 sites) with a survey conducted after wild dog recovery (2011; 48513 trap hours at 97 sites). We placed camera traps in random sites throughout the study area. We used a lag time of 6 minutes between sequential camera images to identify unique camera trap events. We used an exact test with a Poisson distribution to evaluate if detection rates (i.e., images per trap hour) of leopards and jackals had changed between these two periods. Following wild dog recovery, an increase in the number of leopard or jackal detections could obfuscate an effect of predation by wild dogs per se on dik-dik abundance.

Second, to evaluate the potential influence of rainfall on the dik-dik population, we first calculated the cumulative rainfall (mm) over a 6-month period preceding each dik-dik population survey (which represents the average inter-birth period; Kingswood and Kumamoto 1996). We regressed rainfall against r using a GLS analysis, and compared this with models involving the estimated consumption of dik-dik by wild dogs. We also compared the total rainfall per month before and after wild dog recovery using a GLS. A significant, positive effect of rainfall on the population growth rate of dik-dik, combined with an overall decrease in rainfall after wild dog recovery, confounds any negative effect of predation by wild dogs on dik-dik abundance.

Third, we evaluated if dik-dik were more difficult to detect following wild dog recovery. Dik-dik are thought to rely on crypsis to evade predators (Estes 1991, Brashares et al. 2000). If wild dogs reduced the conspicuousness of dik-dik, this could create the perception of reduced abundance. Under this scenario, and following wild dog recovery, the detection distance (i.e., the effective strip width based on distance sampling methodology) should decrease as dik-dik become less conspicuous.

#### Prediction 2: Dik-dik suppress tree abundance

We assessed the effect of dik-dik browsing on three abundant species of tree: Acacia etbaica, Acacia mellifera, and Grewia spp., which comprised approximately 40%, 5-10%, and 8% of tree cover in our study area, respectively (Young et al. 1995). These species are present among all experimental treatments in the before and after wild dog exclusion experiments (see Prediction 3). We also measured the response of the aggregate tree community to dik-dik browsing by pooling the abundances of all tree species (32 species). We measured the effect of browsing by dik-dik per se on tree abundance, using replicated ungulate exclusions that are part of the UHURU (ungulate herbivory under rainfall uncertainty) experiment (Goheen et al. 2013). The UHURU experiment was initiated in 2009 and consists of 36 1-ha fenced areas distributed among three sites that are spread across a spatial gradient in rainfall (Goheen et al. 2013). At each site, there are three 4-ha blocks each consisting of 1-ha treatments that exclude (1) all ungulates (TOTAL); (2) all ungulates  $\geq 40$  kg and  $\geq$ 1.2 m tall, thereby allowing dik-dik (MESO); (3) elephant and giraffes (MEGA); (4) no ungulates (OPEN). Within each 1-ha treatment, we recorded the number of woody plants in the 1.0-2.0 m height class in 2009 and in 2012. We did not include the northern and most arid plots from the UHURU experiment to

maintain consistency with the study area from the before wild dog exclosure experiment (see *Prediction 3*).

To analyze the effect of dik-dik on tree abundance, we calculated the net difference in density of trees in the 1.0-2.0 m height class (individuals· $100 \text{ m}^{-2} \cdot \text{yr}^{-1}$ ) between 2009 and 2012 as the response variable, with treatment (i.e., MESO vs. TOTAL) as the predictor variable, and used a GLS analysis. We then ran a Fisher's combined probability test with a weighted-Z approach for the three species-level GLS models (Whitlock 2005). If dik-dik exerted top-down control on trees, then we expected to see a greater increase in stem density in TOTAL plots (i.e., excludes all ungulates) relative to MESO plots (i.e., those accessible to dik-dik, but not larger than dik-dik).

#### Prediction 3: The effect of dik-dik on tree abundance is reduced in the presence of wild dogs

While Prediction 2 addresses whether dik-dik in isolation have the potential to suppress *A. etbaica*, *A. mellifera*, *Grewia* spp., or the aggregate tree community over a three-year period, Prediction 3 addresses the effect of browsing by all ungulates before and after wild-dog recovery. If wild dog recovery alters the plant community via suppression of dik-dik, then a reduction in browsing by dik-dik should be evident in the presence of other ungulates.

We first measured the effect of ungulate exclusion on tree abundance in 1999–2002, just prior to wild dog recovery (Augustine and McNaughton 2004). This exclusion experiment consisted of three 0.5-ha electrified fenced areas that excluded all ungulates (TOTAL plots), and were paired with 0.5-ha unfenced control areas (OPEN plots). To quantify the effect of ungulate exclusion on tree abundance after wild dog recovery, we compared the TOTAL and OPEN plots from the UHURU experiment (2009–2012).

The trophic cascade hypothesis predicts that differences in tree abundance between OPEN and TOTAL plots should be greater in the before wild dog exclusion experiment than in the after wild dog exclusion experiment; i.e., all else equal, browsing pressure should be reduced in the presence of wild dogs. We tested this prediction using a GLS analysis, with the net difference in the density of trees in the 1.0-2.0 m height class (individuals  $\cdot 100 \text{ m}^{-2} \cdot \text{yr}^{-1}$ ) as the response variable, and an interaction between treatment (OPEN vs. TOTAL plots) and the status of wild dog recovery (before vs. after; hereafter "recovery status") as predictor variables. We included a structured variance term to stabilize heteroscedasticity in residuals. We conducted separate analyses for A. etbaica, A. mellifera, Grewia spp., and the tree community in aggregate, each fit using maximum likelihood to facilitate model selection and selected the best-fitting model using AIC<sub>c</sub>. If the bestfitting model(s) include the interaction term between treatment and recovery status, this may (depending on the direction of the interaction) indicate that wild dogs

suppressed the effect of dik-dik on tree abundance. Thus, if models containing the interaction term had a  $\Delta AIC_c < 2.0$ , we proceeded to refit the model using restricted maximum likelihood estimation, and compared pairwise differences for each combination of treatment and recovery status using a Tukey's honestly significant difference test. In addition, we compared the mean difference in tree abundance (d) for each ungulate exclusion experiment ( $d_s = \text{TOTAL}_s - \text{OPEN}_s$ ) where s indicates recovery status (before vs. after wild dog recovery) at the time of the experiment. We used a pooled standard error to quantify uncertainty in grouped means (Quinn and Keough 2002). The trophic-cascade hypothesis predicts  $d_{\text{before}} > d_{\text{after}}$ ; however, if mean differences are similar, it indicates that the effect of herbivory has not changed appreciably following the recovery of wild dogs.

#### Other potential drivers of tree abundance

In addition to the indirect effect of wild dogs, we considered two alternative drivers of tree abundance. First, we evaluated whether the abundance of browsers other than dik-dik (e.g., impala, giraffe, and elephants) had changed along with wild dog recovery. We compared the energetic demand of all non-dik-dik browsers before (2000-2002) and after wild dog recovery (2008-2011). Population densities of these browsers were quantified while performing the dik-dik population surveys in 2000-2002 (Augustine 2010) and 2008-2011 (T. G. O'Brien and M. F. Kinnaird, unpublished data; see Appendix B: Table B3). Biomass was estimated using the mean adult body size of each browser multiplied by species density. To estimate energetic demand of all browsers, we calculated the mass-specific field metabolic rates as FMR =  $\sum_{j=1}^{n} 4.82 M_j^{0.734}$ , where *M* is the mean biomass density (g/  $km^{-2}$ ) of species *j* and FMR is the energetic demand in kJ/d (Ernest and Brown 2001, Nagy 2005). If the biomass density or energetic demand of browsers (besides dik-dik) has increased with wild dog recovery, it may negate any indirect effect of wild dogs on tree abundance. Likewise, if the density of browsers (besides dik-dik) had decreased with wild dog recovery, it would confound our ability to ascribe increased tree abundance

to the suppression of dik-dik alone. We also considered the possibility that rainfall covaried with the recovery status of wild dogs. Higher rainfall after wild dog recovery could enhance tree survival, growth, and reproduction, and thus confound the indirect effect of wild dogs on tree abundance. Our methods for analyzing rainfall are described in *Prediction 1: Wild dogs suppress dik-dik abundance*.

#### RESULTS

#### Prediction 1: Wild dogs suppress dik-dik abundance

The biomass density of wild dogs on Mpala Research Center peaked between June 2007 and January 2008 at 3938 kg·d<sup>-1</sup>·km<sup>-2</sup>, with a mean biomass density of 1600  $\pm$  266 kg· d<sup>-1</sup>·km<sup>-2</sup> (mean  $\pm$  SE) since recovery in 2002. Dik-dik density was  $145 \pm 4$  individuals/km<sup>2</sup> before wild dog recovery (1999–2002) and 97  $\pm$  7 individuals/km<sup>2</sup> since 2008, corresponding to a  $\sim$ 33% decline in dik-dik abundance ( $F_{1.14} = 27.9$ , P < 0.001; Fig. 1a). The bestfitting model for r (the population growth rate of dikdik) consisted only of the main effect for the energetic demand of wild dogs ( $\beta = -0.78 \pm 0.19$ ,  $F_{1, 13} = 16.0$ , P = 0.002), which far outperformed the next best-fitting model ( $\Delta AIC_c > 6$ ; Appendix B: Table B1).The recruitment index declined by 41%, from 0.17  $\pm$  0.02 (1999-2002) to  $0.10 \pm 0.01$  (2008-2013), and decreased with increasing energetic demand of wild dogs (Fig. 1b). Relative to the pre-denning period, encounter rates with dik-dik decreased by 42% in the denning home range while the den was active (Fig. 1c; Poisson rate parameter,  $\lambda$  [95% CI] = 0.525 [0.36–0.74], P < 0.001). The encounter rate in the control area did not change over the same period of time ( $\lambda = 1.01$  [0.82–1.25], P =0.916]). Two months after den abandonment by wild dogs, the proportionate difference in encounter rates compared to their respective pre-denning period was similar near the den (22%) and in the control area (19%); Fig. 1c). During the active denning phase, the recruitment index declined by 20% within the denning home range, while there was a fivefold increase in the recruitment index over the same period in the control area (Fig. 1d;  $\chi^2 = 5.75$ , P = 0.008). Thus, over both expansive (82 km<sup>2</sup>, 14 years) and localized (31 km, 33 days) spatiotemporal scales, the energetic demand of wild dogs was correlated negatively with abundance and recruitment of dik-dik.

#### Other potential drivers of dik-dik abundance

The decline in the dik-dik population following wild dog recovery could not be explained by an increase in the abundance of other predators, lower rainfall (resource availability), or reduced detectability of dikdik. Compared to before wild dog recovery, the relative abundance of carnivores most likely to consume dik-dik was either the same (leopard,  $\lambda = 1.13$  [0.265, 10.260],  $P \ge 0.999$ ) or significantly less (black-backed jackal,  $\lambda =$ 0.13 [0.041,0.419], P < 0.001) following wild dog recovery.

The observed decline in dik-dik abundance was likely not caused by declining resource availability. On average, 23% more monthly rainfall occurred after wild dog recovery (58.3 ± 4.5 mm, 2003–2013) compared to before wild dog recovery (47.3 ± 6.2 mm, 1999–2002), but this difference was not statistically significant ( $t_{2, 187}$ = 1.12, P = 0.264; Appendix B: Fig. B2). Moreover, we did not find support for an effect of rainfall on population growth of dik-dik (Appendix B: Table B1).

We did not observe a change in the effective strip width of dik-dik during population surveys conducted before (22.7  $\pm$  0.4 m) and after wild dog recovery (24.4  $\pm$  0.5 m). Thus, it is unlikely that the decline in the



FIG. 1. Changes in dik-dik abundance over 14 years in an 82-km<sup>2</sup> area, shown as (a) dik-dik density (black) and the estimated number of dik-dik eaten by wild dogs between dik-dik population surveys (gray). Consumption of dik-dik accounts for energetic content of dik-dik, and the diet composition, total biomass, and days of occupancy by wild dogs in our study area. Error bars show 95% CI. (b) The estimated number of dik-dik eaten by wild dogs was negatively correlated with the recruitment index of dik-dik ( $F_{1,14} = 9.75$ , P = 0.008). Over a finer spatiotemporal scale (34 km, 33 days), suppression of dik-dik by wild dogs was evident on (c) the proportionate difference in encounter rates (dik-dik/km) from the pre-denning period, which decreased by 42% in the denning home range but increased by 10% in a nearby control area where wild dogs foraged much less frequently during the same period (Appendix B: Fig. B1); and on (d) the recruitment index, which decreased near the den but increased in the control area during the same time. Responses during the active- and post-denning periods are equal to the pre-denning period at a value of 0.

abundance of dik-dik was an artifact of heightened crypsis following wild dog recovery.

#### Prediction 2: Dik-dik suppress tree abundance

By themselves, dik-dik significantly reduced the abundance of *A. mellifera* but their effect on the abundance of *A. etbaica*, *Grewia* spp., and the aggregate tree community was not statistically detectable (Fig. 2). Over a three-year period, the abundance of *A. mellifera* in TOTAL plots (excluded dik-dik and other ungulates) increased by 84% relative to MESO plots (allowed dik-dik, excluded other ungulates;  $t_{1,10} = 2.88$ , P = 0.016). The abundance of *A. etbaica*, *Grewia* spp., and the aggregate tree community increased by 54% ( $t_{1,10} = 1.37$ , P = 0.201), 52% ( $t_{1,10} = 1.49$ , P = 0.167), and 40% ( $t_{1,10} = 0.96$ , P = 0.360), respectively, in

TOTAL plots relative to MESO plots (Fig. 2). This result indicates that dik-dik had a tendency to exert top-down control on these species. A Fisher's combined probability test (Whitlock 2005) of *A. mellifera*, *A. etbaica*, and *Grewia* spp. indicated that this collective difference was statistically significant (P = 0.011). Thus, between 2009 and 2012, dik-dik alone contributed to the suppression of three tree species and the aggregate tree community, with *A. mellifera* being the most sensitive species to variation in herbivory by dik-dik (Fig. 2).

#### Prediction 3: The effect of dik-dik on tree abundance is reduced in the presence of wild dogs

Generally, tree abundance increased at a faster rate after wild dog recovery than before wild dog recovery.



FIG. 2. Top-down regulation of dik-dik on tree abundance, shown as the net rate of change in tree density between 2009 and 2012 for individuals in the 1.0-2.0 m height class. TOTAL plots excluded dik-dik and all other browsers, while MESO plots permitted access by dik-dik (5 kg), but not larger browsers. Error bars indicate  $\pm$ SE; asterisk indicates significance (P < 0.05) for a generalized least squares analysis.

For example, there was a significant increase in stem abundance for the aggregate tree community in OPEN plots after wild dog recovery compared to OPEN plots before wild dog recovery ( $t_{1,9} = -2.96$ , P = 0.021; Appendix B: Fig. B3). This effect was in the same direction, but was not significant for *A. mellifera* ( $t_{1,9} =$ -1.67, P = 0.138), *A. etbaica* ( $t_{1,9} = -2.05$ , P = 0.080), and *Grewia* spp. ( $t_{1,9} = -1.45$ , P = 0.190). A Fisher's combined probability test (Whitlock 2005) of *A. mellifera*, *A. etbaica*, and *Grewia* spp. indicated that this collective difference was statistically significant (P =0.026). Superficially, the positive relationship between the recovery of wild dogs and increased tree abundance in OPEN plots is consistent with the trophic-cascade hypothesis.

Critically, however, the net effect of herbivory, as measured by comparing tree abundance in OPEN vs. TOTAL plots, was not reduced in the presence of wild dogs. None of the best-fitting ( $\Delta AIC_c < 2$ ) models for the effect of ungulate exclusion on any tree species included an interaction of treatment and recovery status (Appendix B: Table B2). By itself, ungulate exclusion was included in the best-fitting models for all three tree species, showing that browsing has an important effect on tree abundance. The status of wild dog recovery was present in the best-fitting models for A. etbaica, A. mellifera, and Grewia spp. (Appendix B: Table B2); however, the direction of this effect was negative, in contrast to the central prediction of the trophic-cascade hypothesis (Fig. 3). These results indicate that the effect of herbivory by the entire community of browsing ungulates is either unchanged or has intensified following wild dog recovery: in other words, we did not find evidence for a trophic cascade.

#### Other potential drivers of tree abundance

A slight increase in the abundance of other browsers and variation in rainfall may explain why the effect of herbivory was not reduced by wild dog recovery. The total biomass density of browsing ungulates (excluding dik-dik) increased slightly from 4129 kg/km<sup>2</sup> before wild dog recovery to 4178 kg/km<sup>2</sup> after wild dog recovery (Appendix B: Table B3), and the energetic demand of non-dik-dik browsers increased slightly by 7%, from 450 MJ/km<sup>2</sup> to 480 MJ/km<sup>2</sup>, following wild dog recovery. In addition, there was a statistically nonsignificant 36% increase in mean monthly rainfall from before (44.8 ± 6.7 mm [1999–2002]) to after wild dog recovery (60.9 ± 7.6 mm [2009–2012];  $F_{1.94} = 1.19$ , P = 0.278).

#### DISCUSSION

We did not find support for a trophic cascade following the recovery of African wild dogs, in spite of (1) the suppression of the dik-dik population by wild dogs, (2) dik-dik's suppression of tree abundance, and (3) the positive correlation between tree and wild dog abundance. Trophic cascades arise when plant abundance is increased by the alteration of top-down forces through at least two sequential trophic levels. These forces must be demonstrably stronger than other factors that regulate species abundance. We evaluated a number of potentially limiting factors for dik-dik populations and could find no explanation more parsimonious than predation by wild dogs. Consistent with Augustine and McNaughton (2004), we also demonstrated that dik-dik by themselves suppress the abundance of at least some



FIG. 3. The effect of browsing on tree abundance before (1999–2002) and after (2009–2012) recovery of wild dogs, where d = TOTAL - OPEN, i.e., the difference between the mean tree abundance of TOTAL plots (excludes all ungulates) and OPEN (accessible to all ungulates) plots over a three-year period for individuals in the 1.0–2.0 m height class. Tree abundances for each combination of treatment (TOTAL vs. OPEN) and experiment (before vs. after wild dog recovery) are shown in Appendix B: Fig. B3. Error bars indicate ±SE.

tree species. Although top-down forces are strong within this ecosystem, the recovery of wild dogs did not counteract intense browsing pressure. Below, we discuss potential mechanisms that may have prevented us from detecting the cascading effects of wild dogs, and highlight the implications of our findings for the restoration of large carnivores in savanna ecosystems.

A lingering question from our study is why wild dogs, in spite of their demonstrably strong effect on the population of the most abundant browser in this system, did not generate a detectable trophic cascade. One explanation, rooted in food web theory (Yodzis 1988), suggests that indirect effects take longer to manifest than direct effects (but see Menge 1997). In our study, the 20-year absence of wild dogs prior to their recovery, coupled with the seven-year difference between the initiation of our two herbivore exclusion experiments, may not have been long enough to detect a change in the abundance of long-lived plants like Acacia spp., Grewia spp., and other trees in this community. A second explanation is that the 36% increase in mean monthly rainfall following the recovery of wild dogs may have overridden any signal of reduced browsing by dik-dik. Across African savannas, tree abundance is limited by rainfall below a mean annual precipitation (MAP) of 650 mm, and more so by herbivores and fire above a MAP of 650 mm (Sankaran et al. 2005). During the before-recovery exclosure experiment, MAP was below this threshold  $(537 \pm 100 \text{ mm})$  but exceeded it following wild dog recovery (730  $\pm$  146 mm). Because of increased rainfall, and in spite of declining dik-dik abundance, the per capita (i.e., per individual herbivore) effect of browsing on tree abundance may have increased following the recovery of wild dogs (but see Louthan et al. 2013). A third and perhaps more promising explanation is that this community of browsing ungulates constitutes a reticulate food web that dampened the strength of indirect effects resulting from wild dog recovery (Polis and Strong 1996), such that forage that would have otherwise been released by the suppression of dik-dik may have been consumed by other species.

Correlative studies and natural experimentation often are the only means through which to investigate trophic cascades at scales commensurate with the movements and lifespans of large mammals and their prey. Our results highlight the advantages of incorporating manipulative approaches to explicitly quantify the constituent interactions that create trophic cascades. Trophic cascades require that top-down limitation, by way of at least two direct interactions (e.g., carnivore-herbivore, herbivore-plant), give rise to an indirect interaction (e.g., carnivore-plant). A strong inferential approach to the study of trophic cascades would therefore (1) directly quantify each of the three interactions hypothesized to comprise the cascade (Schmitz 2010) and (2) evaluate alternative explanations that can produce patterns similar to trophic

cascades (Estes et al. 1998, Post et al. 1999, Hebblewhite et al. 2005, Peterson et al. 2014). A number of studies, most of which were conducted in temperate biomes, have interpreted a positive correlation between the biomass of plants and large carnivores as evidence of a trophic cascade (e.g., Ripple et al. 2001, Callan et al. 2013, Kuijper et al. 2013). This approach addresses patterns consistent with indirect effects, but quantifies neither the response of herbivores to carnivores, nor the response of plants to herbivores. Consequently, and because of the absence of a demonstrated mechanism, some of these studies have generated lively debate centered on the standards of evidence needed to demonstrate causation in the trophic-cascade hypothesis (Mech 2012, Winnie 2012, 2014, Allen et al. 2013, Kauffman et al. 2013, Beschta et al. 2014). In the few studies to manipulate predation by large carnivores through removal experiments, trophic cascades did not occur even though carnivore-herbivore and herbivoreplant interactions were strong (Sinclair et al. 2000, Maron and Pearson 2011). In our study, wild dogs suppressed dik-dik, dik-dik suppressed trees, and tree abundance outside of herbivore exclusion plots increased following wild dog recovery, a pattern that would otherwise suggest a trophic cascade. However, counter to the predictions of the trophic-cascade hypothesis, we found that the effect of herbivory on tree abundance was not reduced following wild dog recovery. Consequently, the claim that trophic cascades are a universal property of ecosystems (sensu Terborgh et al. 2010) may be premature without evidence from a greater number of ecological communities, combined with more concerted efforts to evaluate alternative hypotheses, especially in systems where experiments often are not used to test predictions.

Across savannas, tree cover is a key determinant of both ecosystem dynamics and rural livelihoods, as it affects nutrient cycles (Belsky 1994, Treydte et al. 2007), surface water retention (Scholes and Archer 1997, Smit and Rethman 2000), forage for both wild and domestic herbivores (Odadi et al. 2009, 2011), and household fuel availability in many areas (Chambers and Longhurst 1986). Factors influencing tree cover are therefore an important consideration for ecological and conservation-related research. Rainfall, soils, natural disturbance, and herbivory are widely recognized drivers of tree cover in African savannas (Sankaran et al. 2005, Bond 2008, Lehmann et al. 2014). If large carnivores also contribute to the regulation of tree cover via trophic cascades, it could have profound impacts on the livelihoods of people and ecosystem function (Estes et al. 2011, Ripple et al. 2014). Although indirect effects are an important process in some food webs (Terborgh et al. 2010), further study is needed to understand the ecological contexts in which the restoration of large carnivores will trigger trophic cascades in African savannas.

#### ACKNOWLEDGMENTS

This research was supported by grants from the Natural Sciences and Engineering Research Council (A. T. Ford, J. R. Goheen), a Killam Pre-doctoral Fellowship (A. T. Ford), the American Society of Mammalogists (A. T. Ford), University of British Columbia (A. T. Ford), University of Wyoming (J. R. Goheen), Mpala Research Trust (M. F. Kinnaird), USAID (M. F. Kinnaird, T. G. O'Brien), Wildlife Conservation Society (M. F. Kinnaird, T. G. O'Brien, R. Woodroffe), National Geographic Society (R. Woodroffe), African Wildlife Foundation (R. Woodroffe), and Denver Zoo (R. Woodroffe).

#### LITERATURE CITED

- Allen, B. L., P. J. S. Fleming, L. R. Allen, R. M. Engeman, G. Ballard, and L. K. P. Leung. 2013. As clear as mud: a critical review of evidence for the ecological roles of Australian dingoes. Biological Conservation 159:158–174.
- Augustine, D. J. 2010. Response of native ungulates to drought in semi-arid Kenyan rangeland. African Journal of Ecology 48:1009–1020.
- Augustine, D. J., and S. J. McNaughton. 2004. Regulation of shrub dynamics by native browsing ungulates on East African rangeland. Journal of Applied Ecology 41:45–58.
- Belsky, A. J. 1994. Influences of trees on savanna productivity: tests of shade, nutrients, and tree–grass competition. Ecology 75:922–932.
- Beschta, R. L., C. Eisenberg, J. W. Laundré, W. J. Ripple, and T. P. Rooney. 2014. Predation risk, elk, and aspen: comment. Ecology 95:2669–2671.
- Beschta, R. L., and W. J. Ripple. 2012. The role of large predators in maintaining riparian plant communities and river morphology. Geomorphology 157:88–98.
- Beschta, R. L., and W. J. Ripple. 2013. Are wolves saving Yellowstone's aspen? A landscape-level test of a behaviorally mediated trophic cascade: comment. Ecology 94:1420–1425.
- Bond, W. J. 2008. What limits trees in C<sub>4</sub> grasslands and savannas? Annual Review of Ecology Evolution and Systematics 39:641–659.
- Brashares, J. S., T. Garland, and P. Arcese. 2000. Phylogenetic analysis of coadaptation in behavior, diet, and body size in the African antelope. Behavioral Ecology 11:452–463.
- Callan, R., N. P. Nibbelink, T. P. Rooney, J. E. Wiedenhoeft, and A. P. Wydeven. 2013. Recolonizing wolves trigger a trophic cascade in Wisconsin (USA). Journal of Ecology 101: 837–845.
- Chambers, R., and R. Longhurst. 1986. Trees, seasons and the poor. IDS Bulletin 17:44–50.
- Croll, D. A., J. L. Maron, J. A. Estes, E. M. Danner, and G. V. Byrd. 2005. Introduced predators transform subarctic islands from grassland to tundra. Science 307:1959–1961.
- Ernest, M., and J. H. Brown. 2001. Homeostasis and compensation: the role of species and resources in ecosystem stability. Ecology 82:2118–2132.
- Estes, J. A., et al. 2011. Trophic Downgrading of Planet Earth. Science 333:301–306.
- Estes, J. A., M. T. Tinker, T. M. Williams, and D. F. Doak. 1998. Killer whale predation on sea otters linking oceanic and nearshore ecosystems. Science 282:473–476.
- Estes, R. D. 1991. The behavior guide to African mammals: including hoofed mammals, carnivores, primates. University of California Press, Oakland, California, USA.
- Finke, D. L., and R. F. Denno. 2004. Predator diversity dampens trophic cascades. Nature 429:407–410.
- Ford, A. T., J. R. Goheen, T. O. Otieno, L. Bidner, L. A. Isbell, T. M. Palmer, D. Ward, R. Woodroffe, and R. M. Pringle. 2014. Large carnivores make savanna tree communities less thorny. Science 346:346–349.
- Goheen, J. R., T. M. Palmer, G. K. Charles, K. M. Helgen, S. N. Kinyua, J. E. Maclean, B. L. Turner, H. S. Young, and R. M. Pringle. 2013. Piecewise disassembly of a large-

herbivore community across a rainfall gradient: the UHURU experiment. PLoS ONE 8:e55192.

- Hairston, N. G., F. E. Smith, and L. B. Slobodkin. 1960. Community structure, population control and competition. American Naturalist 94:421–425.
- Hebblewhite, M., C. A. White, C. G. Nietvelt, J. A. McKenzie, T. E. Hurd, J. M. Fryxell, S. E. Bayley, and P. C. Paquet. 2005. Human activity mediates a trophic cascade caused by wolves. Ecology 86:2135–2144.
- Holling, C. S. 1959. Some characteristics of simple types of predation and parasitism. Canadian Entomologist 91:385– 398.
- Hopcraft, J. G. C., H. Olff, and A. R. E. Sinclair. 2010. Herbivores, resources and risks: alternating regulation along primary environmental gradients in savannas. Trends in Ecology & Evolution 25:119–128.
- Kauffman, M. J., J. F. Brodie, and E. S. Jules. 2010. Are wolves saving Yellowstone's aspen? A landscape-level test of a behaviorally mediated trophic cascade. Ecology 91:2742– 2755.
- Kauffman, M. J., J. F. Brodie, and E. S. Jules. 2013. Are wolves saving Yellowstone's aspen? A landscape-level test of a behaviorally mediated trophic cascade: reply. Ecology 94: 1425–1431.
- Kingswood, S. C., and A. T. Kumamoto. 1996. Madoqua guentheri. Mammalian Species 539:1–10.
- Komers, P. E. 1996. Obligate monogamy without paternal care in Kirk's dikdik. Animal Behaviour 51:131–140.
- Kuijper, D. P. J., C. de Kleine, M. Churski, P. van Hooft, J. Bubnicki, and B. Jedrzejewska. 2013. Landscape of fear in Europe: wolves affect spatial patterns of ungulate browsing in Bialowieza Primeval Forest, Poland. Ecography 36:1263– 1275.
- Kuker, K., and L. Barrett-Lennard. 2010. A re-evaluation of the role of killer whales *Orcinus orca* in a population decline of sea otters *Enhydra lutris* in the Aleutian Islands and a review of alternative hypotheses. Mammal Review 40:103– 124.
- Lehmann, C. E. R., et al. 2014. Savanna vegetation-fire-climate relationships differ among continents. Science 343:548–552.
- Leibold, M. A. 1989. Resource edibility and the effects of predators and productivity on the outcome of trophic interactions. American Naturalist 134:922–949.
- Louthan, A. M., D. F. Doak, J. R. Goheen, T. M. Palmer, and R. M. Pringle. 2013. Climatic stress mediates the impacts of herbivory on plant population structure and components of individual fitness. Journal of Ecology 101:1074–1083.
- Maron, J. L., and D. E. Pearson. 2011. Vertebrate predators have minimal cascading effects on plant production or seed predation in an intact grassland ecosystem. Ecology Letters 14:661–669.
- Mech, L. D. 2012. Is science in danger of sanctifying the wolf? Biological Conservation 150:143–149.
- Menge, B. A. 1997. Detection of direct versus indirect effects: Were experiments long enough? American Naturalist 149: 801–823.
- Mooney, K. A., R. Halitschke, A. Kessler, and A. A. Agrawal. 2010. Evolutionary trade-offs in plants mediate the strength of trophic cascades. Science 327:1642–1644.
- Nagy, K. A. 2005. Field metabolic rate and body size. Journal of Experimental Biology 208:1621–1625.
- Newsome, T. M., G. A. Ballard, C. R. Dickman, P. J. S. Fleming, and R. van de Ven. 2013. Home range, activity and sociality of a top predator, the dingo: a test of the Resource Dispersion Hypothesis. Ecography 36:914–925.
- Odadi, W. O., M. K. Karachi, S. A. Abdulrazak, and T. P. Young. 2011. African wild ungulates compete with or facilitate cattle depending on season. Science 333:1753–1755.
- Odadi, W. O., J. B. Okeyo-Owuor, and T. P. Young. 2009. Behavioural responses of cattle to shared foraging with wild

- Peterson, R. O., J. A. Vucetich, J. M. Bump, and D. W. Smith. 2014. Trophic cascades in a multicausal world: Isle Royale and Yellowstone. Annual Review of Ecology, Evolution, and Systematics 45:325–345.
- Polis, G. A., A. L. W. Sears, G. R. Huxel, D. R. Strong, and J. Maron. 2000. When is a trophic cascade a trophic cascade? Trends in Ecology & Evolution 15:473–475.
- Polis, G. A., and D. R. Strong. 1996. Food web complexity and community dynamics. American Naturalist 147:813–846.
- Post, E., R. O. Peterson, N. C. Stenseth, and B. E. McLaren. 1999. Ecosystem consequences of wolf behavioural response to climate. Nature 401:905–907.
- Quinn, G. P., and M. J. Keough. 2002. Experimental design and data analysis for biologists. Cambridge University Press, Cambridge, UK.
- Ripple, W. J., et al. 2014. Status and ecological effects of the world's largest carnivores. Science 343.
- Ripple, W. J., E. J. Larsen, R. A. Renkin, and D. W. Smith. 2001. Trophic cascades among wolves, elk and aspen on Yellowstone National Park's northern range. Biological Conservation 102:227–234.
- Sankaran, M., et al. 2005. Determinants of woody cover in African savannas. Nature 438:846–849.
- Schmitz, O. 2010. Resolving ecosystem complexity. Princeton University Press, Princeton, New Jersey, USA.
- Schmitz, O. J. 1994. Resource edibility and trophic exploitation in an old-field food-web. Proceedings of the National Academy of Sciences USA 91:5364–5367.
- Schmitz, O. J., V. Krivan, and O. Ovadia. 2004. Trophic cascades: the primacy of trait-mediated indirect interactions. Ecology Letters 7:153–163.
- Scholes, R., and S. Archer. 1997. Tree-grass interactions in savannas 1. Annual Review of Ecology and Systematics 28: 517–544.
- Sinclair, A. R. E., and P. Arcese. 1995. Population consequences of predation-sensitive foraging: the Serengeti wildebeest. Ecology 76:882–891.
- Sinclair, A. R. E., C. J. Krebs, J. M. Fryxell, R. Turkington, S. Boutin, R. Boonstra, P. Seccombe-Hett, P. Lundberg, and L. Oksanen. 2000. Testing hypotheses of trophic level interactions: a boreal forest ecosystem. Oikos 89:313–328.
- Smit, G., and N. Rethman. 2000. The influence of tree thinning on the soil water in a semi-arid savanna of southern Africa. Journal of Arid Environments 44:41–59.

- Strong, D. R. 1992. Are trophic cascades all wet? Differentiation and donor-control in speciose ecosystems. Ecology 73: 747–754.
- Terborgh, J., R. D. Holt, and J. A. Estes. 2010. Trophic cascades: what they are, how they work, and why they matter. Pages 1–18 *in* J. Terborgh and J. A. Estes, editors. Trophic cascades: predators, prey, and the changing dynamics of nature. Island Press, Washington, D.C., USA.
- Thibault, K. M., S. K. M. Ernest, and J. H. Brown. 2010. Redundant or complementary? Impact of a colonizing species on community structure and function. Oikos 119: 1719–1726.
- Treydte, A. C., I. Heitkönig, H. H. Prins, and F. Ludwig. 2007. Trees improve grass quality for herbivores in African savannas. Perspectives in Plant Ecology, Evolution and Systematics 8:197–205.
- Tschanz, B., L. F. Bersier, and S. Bacher. 2007. Functional responses: a question of alternative prey and predator density. Ecology 88:1300–1308.
- van der Stap, I., M. Vos, A. M. Verschoor, N. R. Helmsing, and W. M. Mooij. 2007. Induced defenses in herbivores and plants differentially modulate a trophic cascade. Ecology 88: 2474–2481.
- Whitlock, M. 2005. Combining probability from independent tests: the weighted Z-method is superior to Fisher's approach. Journal of Evolutionary Biology 18:1368–1373.
- Winnie, J. 2014. Predation risk, elk, and aspen: reply. Ecology 95:2671–2674.
- Winnie, J. A. 2012. Predation risk, elk, and aspen: tests of a behaviorally mediated trophic cascade in the Greater Yellowstone Ecosystem. Ecology 93:2600–2614.
- Woodroffe, R. 2011. Demography of a recovering African wild dog (*Lycaon pictus*) population. Journal of Mammalogy 92: 305–315.
- Woodroffe, R., P. A. Lindsey, S. S. Romanach, and S. Ranah. 2007. African wild dogs (*Lycaon pictus*) can subsist on small prey: implications for conservation. Journal of Mammalogy 88:181–193.
- Yodzis, P. 1988. The indeterminacy of ecological interactions as perceived through perturbation experiments. Ecology 69: 508–515.
- Young, T. P., N. Patridge, and A. Macrae. 1995. Long-term glades in *Acacia* bushland and their edge effects in Laikipia, Kenya. Ecological Applications 5:97–108.
- Zuur, A., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed effects models and extensions in ecology with R. Springer, New York, New York, USA.

SUPPLEMENTAL MATERIAL

#### **Ecological Archives**

Appendices A and B are available online: http://dx.doi.org/10.1890/14-2056.1.sm

## **Ecological Archives E096-238-A1**

## Adam T. Ford, Jacob R. Goheen, David J. Augustine, Margaret F. Kinnaird, Timothy G. O'brien, Todd M. Palmer, Robert M. Pringle, and Rosie Woodroffe. 2015. Recovery of African wild dogs suppresses prey but does not trigger a trophic cascade. *Ecology* 96:2705–2714. <u>http://dx.doi.org/10.1890/14-2056.1</u>

APPENDIX A. Description of study area, methods for assessing the abundance of dik-dik and wild dogs, hypotheses tested, and statistical methods used.

## Study area

We undertook our research at the Mpala Research Centre (MRC), a private wildlife conservancy in Laikipia County, Kenya (0° 17' N, 37° 53' E). This semi-arid savanna is characterized by a discontinuous overstory comprising about 28% of total cover and dominated by *Acacia brevispica*, *A. etbaica*, *A. mellifera*, *Grewia* sp., and *Croton* sp.(Young et al. 1995). MRC has a mean annual rainfall of 508 mm, varying steeply along a north-south gradient (Goheen et al. 2013) with marked inter-annual variation (Augustine 2010).

## Recovery of wild dogs

Estimated biomass of wild dogs on MRC was derived from an ongoing study of > 30 packs, distributed over 12,000 km<sup>2</sup> (<u>Woodroffe 2011</u>). Using GPS and VHF telemetry, we quantified the movement patterns, denning periods, and composition of 9 packs at MRC since 2002. Collars were fit on 1–2 individuals in each pack. Because packs are highly cohesive, we interpreted the telemetry data from collared individuals as representative of the entire pack. To quantify the biomass density of wild dogs on MRC, we first calculated the composition of adults (ca. 23 kg) and juveniles (5-22 kg) in each pack, and summed the estimated mass of these individuals. The biomass estimate of individual juveniles was adjusted over time as they matured (<u>Woodroffe 2011</u>). We used telemetry to estimate the number of days that each pack spent at MRC, yielding an estimate of wild dog biomass days (in kg·days) over the study area. We performed this calculation for the 6 months preceding each population survey for dik-dik (see below) up to January 2013, as well as in January and June of the years where surveys for dik-diks were not conducted (2003–2007). Using GPS telemetry and field observations, we identified a denning event from a single wild-dog pack comprised of 19 adults and 12 pups in 2011, during which time individuals foraged intensively within 3 km of their den (Appendix B, Fig. B1).

## Monitoring the abundance of dik-dik

We monitored dik-dik abundance using line-transect sampling methods. In a previous study, one of us (DJA) performed linetransect sampling (Buckland et al. 2007) to estimate dik-dik population density along four transects (1.5–3.3 km each) in 2000 and six transects (1.5–3.3 km each) in 2001–2002 (Augustine 2010). In each survey, transects were each driven four times in 2000 and six times in 2001–2002, for a total effort of 35.9 km to 92.8 km per survey. A total of six surveys (3 in 2000, 2 in 2001, 1 in 2002) were performed prior to wild dog recovery. Beginning in June 2008 (i.e., 6 years after wild dogs had recolonized MRC) we monitored the same study area as Augustine (2010) during semi-annual surveys. Each of these surveys consisted of 20 ca. 2 km transects that were sampled once per survey, for a total effort of ca. 40 km per survey. A total of 12 surveys between June 2008 and January 2014.

Line transect sampling was always performed with two observers and one driver, from a vehicle travelling at 10 km·h<sup>-1</sup>. Distances to animals were estimated using a laser rangefinder and bearings were estimated using a handheld compass. Density estimates and confidence intervals were calculated in Program DISTANCE using a hazard function and cosine series expansion, with observations filtered to a maximum of 72 m from the transects. We filtered data to meet the assumptions of the distance sampling (Buckland et al. 2005), resulting in the removal of 5% of dik-dik observations that were found at the farthest distances from the road.

TABLE A1. Summary of hypotheses, predictions, and data used to evaluate the cascading effects of wild dog recovery in Laikipia, Kenya.

Hypothesis	Description	Prediction if trophic cascade hypothesis is correct	Response variable	Response variable methods or data used	Predictor variable	Predictor variable methods or data used	Statistical model*
		1	1				

H1 - main	Wild dogs suppress dik- dik	Dik-dik density is lower following wild dog recovery	Dik-dik density pre- and post-wild dog recovery	Distance sampling surveys (1999– 2002; 2008– 2013)	Wild dog recovery status	Telemetry and field observations of pack biomass and occupancy	1
	Wild dogs suppress dik- dik	Wild dog biomass has a negative effect on the population growth rate of dik-dik	Population growth rate of dik-dik	Derived from estimates of dik- dik population density pre- and post-wild dog recovery (1999– 2002; 2008– 2013)	Wild dog energetic demand (2008– 2013)	Telemetry and field observations of pack biomass and occupancy (2008– 2013); diet composition of wild dogs; energy available in a dik- dik	2
	Wild dogs suppress dik- dik	Wild dog biomass has a negative effect on dik-dik recruitment	Recruitment index	Derived from group sizes during distance sampling surveys (1999– 2002; 2008– 2013)	Wild dog energetic demand (2008– 2013)	Telemetry and field observations of pack biomass and occupancy (2008– 2013); diet composition of wild dogs; energy available in a dik- dik	3
	Wild dogs suppress dik- dik	Proximity to a wild dog den decreases dik- dik abundance	Encounter rate of dik-dik before, during, and after denning	Encounter rates along road transects within and adjacent to the denning home range of wild dogs (2011–2012)	Before vs. after wild dog denning in areas near and away from the active den site	Telemetry and field observations of wild dog movements	4
	Wild dogs suppress dik- dik	Proximity to a wild dog den has a negative effect on dik-dik recruitment	Recruitment index	Encounter rates along road transects within and adjacent to the denning home range of wild dogs (2011–2012)	Before vs after wild dog denning in areas near and away from the active den site	Telemetry and field observations of wild dog movements	5
H1- alternatives	Predators other than wild dogs suppress dik- dik	Equal or less abundance of jackals and leopards post- wild dog recovery	Relative abundance (detections per unit effort)	Camera trapping pre- (2000– 2002; 7364 trap hours at 19 sites) and post-wild dog (2011; 48513 trap hours at 97 sites)	Wild dog recovery status	Telemetry and field observations of pack biomass and occupancy	6
	Rainfall limits dik-dik abundance	Equal or greater rainfall following the	Population growth rate of dik-dik	Derived from estimates of dik- dik population	Total rainfall in the 6 months preceding the	Rainfall gauge	7

		recovery of wild dogs		density pre- and post-wild dog recovery (1999– 2002; 2008– 2013).	dik-dik density surveys (1999– 2002; 2008– 2013)		
	Dik-dik are more difficult to detect because of wild dogs	Detection distance is the same following wild dog recovery	Detection distance	Derived from distance sampling surveys (1999– 2002; 2008– 2013)	Wild dog recovery status	Telemetry and field observations of pack biomass and occupancy	8
H2 - main	Dik-dik suppress trees	Trees will be more abundant in the absence of dik-dik than in the presence of dik-dik	Net difference in density of trees in the 1.0–2.0 m height class (individuals·100 m <sup>-2</sup> ·yr <sup>-1</sup> )	Tree census in 2009 and 2012	Dik-dik access (MESO vs. TOTAL)	Size-selective herbivore exclosure that either allows dik-dik, but not larger browsers, or excludes dik-dik and larger browsers	9
H3 - main	The effect of dik-dik on tree abundance is reduced in the presence of wild dogs	The net difference in density of trees between exclosures and controls is greater pre-wild dog recovery than post-wild dog recovery	Net difference in density of trees in the 1.0–2.0 m height class (individuals·100 m <sup>-2</sup> ·yr <sup>-1</sup> )	Tree census in 2009–2012, and in 1999–2002.	Ungulate access (OPEN vs. TOTAL)	Exclosures that allow all (OPEN) or no (TOTAL) ungulates	10
H3 - alternatives	Browsers other than dik-dik suppress trees	The density of browsers other than dik-dik is the same or greater following wild dog recovery	Biomass density of browser pre and post-wild dog recovery	Distance sampling surveys (1999– 2002; 2008– 2013)	Wild dog recovery status	Telemetry and field observations of pack biomass and occupancy	NA
	Rainfall limits tree abundance	Rainfall increased following wild dog recovery	Monthly rainfall	Rainfall gauge	Wild dog recovery status	Telemetry and field observations of pack biomass and occupancy	11

\* See Table A2

TABLE A2. Structure of statistical models.

Statistical		NT /
model	Code implementation	Notes
1	gls(DENSITYdd ~ status, weights = varIdent (form =~ status))	
2*	$gls(r_d \sim EATEN, correlation = corARMA (c(0.7, 2), p = 1, q=1),$	
	weights = varExp (form =~ EATEN))	
3	gls ( $RI \sim x$ )	
4	poisson.test (c(Ac, Pr+Po),c(Dn,Dn+Dn))	
	poisson.test (c(Ac, Pr+Po),c(Ct,Ct+Ct))	
5	prop.test (RI <sub>ContolBef</sub> , RI <sub>ControlActive</sub> )	
	prop.test ( <i>RI</i> <sub>DenBef</sub> , <i>RI</i> <sub>DenActive</sub> )	
6	poisson.test (c(Njackal <sub>pre</sub> , Njackal <sub>post</sub> ),(Effort <sub>pre</sub> , Effort <sub>post</sub> ))	
	$poisson.test~(c(Nleopard_{pre},~Nleopard_{post}),(Effort_{pre},~Effort_{post}))$	
7*	$gls(r_d \sim Rain, correlation = corARMA (c(0.7,.2), p = 1, q=1)$	
8	t.test (Dist ~ status)	
9	gls(DENSEDIFF ~ TREATMENTdd, weights = varIdent (form = ~	Separate GLS tests were run for 3
	1  TREATMENTdd))	tree species and 1 for the
		combined tree community
	combine.test (p=(each species specific models for #9 here), method	
	= "z.transform")	
10	gls(DENSEDIFF ~ TREATMENTall + status +	Separate GLS tests were run for 3
	TREATMENTall:status, weights = varIdent (form = ~	tree species and 1 for the
	1 status*TREATMENTall))	combined tree community
11	gls( <i>Rain_month</i> ~ <i>status</i> , correlation = corAR1(),	

### Symbols:

DENSITYdd = dik-dik density (individuals / km<sup>2</sup>) [continuous]

- status = pre- vs post-wild dog recovery [factor]
- rd = population growth rate of dik-dik (individuals/survey) [continuous]
- x = the estimated consumption of dik-dik by wild dogs (individuals/km<sup>2</sup>/survey) [continuous]
- *RI* = recruitment index [continuous]
- Pr, Ac, Po = pre-, active-, and post-wild dog denning, respectfully (number of dik-dik)[count]
- *Bef* = survey period before active denning
- Active = survey period during active denning
- Dn, Ct = transect length denning home range, control area, respectfully (km) [continuous]
- Njackal = number of camera trap detections for jackal [count]
- Nleopard = number of camera trap detections for jackal [count]
- Effort = number of camera trap hours (sites x hours active) [continuous]
- *Pre* = pre-wild dog recovery
- Post = post wild dog recovery
- Rain = total rainfall in the 6 months prior to survey (mm) [continuous]
- Dist = detection distance during each survey (m) [continuous]
- DENSEDIFF = the net difference in density of trees in the 1.0 2.0 m height class
- (individuals·100 m<sup>-2</sup>·yr<sup>-1</sup>) [continuous]
- TREATMENTdd = dik-dik exclosure treatment: MESO -allows dik-dik, but not larger browsers;
- TOTAL excludes dik-dik and larger browser[factor]
- TREATMENTall = ungulate exclosure treatment: OPEN allows all ungulates; TOTAL -
- excludes all ungulates [factor]
- DENSITYbr = density of browsers (individuals / km<sup>2</sup>) [continuous]
- *Rain\_month* = total rainfall in a month (mm) [continuous]
- \* these tests were combined into a model selection process show in Appendix B; Table B1

LITERATURE CITED

Augustine, D. J. 2010. Response of native ungulates to drought in semi-arid Kenyan rangeland. African Journal of Ecology 48:1009–1020.

Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 2005. Distance sampling. Wiley Online Library.

Buckland, S. T., D. R. Anderson, K. P. Burnham, and L. Thomas. 2007. Advanced Distance Sampling: Estimating Abundance of Biological Populations. Oxford University Press, Incorporated.

Goheen, J. R., T. M. Palmer, G. K. Charles, K. M. Helgen, S. N. Kinyua, J. E. Maclean, B. L. Turner, H. S. Young, and R. M. Pringle. 2013. Piecewise disassembly of a large-herbivore community across a rainfall gradient: the UHURU experiment. PLoS ONE 8:e55192.

Woodroffe, R. 2011. Demography of a recovering African wild dog (*Lycaon pictus*) population. Journal of Mammalogy 92:305–315.

Young, T. P., N. Patridge, and A. Macrae. 1995. Long-term glades in Acacia bushland and their edge effects in Laikipia, Kenya. Ecological Applications 5:97–108.

[Back to E096-238]

## **Ecological Archives E096-238-A2**

Adam T. Ford, Jacob R. Goheen, David J. Augustine, Margaret F. Kinnaird, Timothy G. O'brien, Todd M. Palmer, Robert M. Pringle, and Rosie Woodroffe. 2015. Recovery of African wild dogs suppresses prey but does not trigger a trophic cascade. *Ecology* 96:2705–2714. http://dx.doi.org/10.1890/14-2056.1

APPENDIX B. Supplementary results showing the distribution of wild dogs before and after denning, rainfall since 1998, change in stem density for 4 groups of tree species in experiments established before and after wild dog recovery, model selection results for factors affecting dik-dik abundance, and the abundance of dominant ungulate species before and after wild dog recovery.



FIG. B1. Location of transects near and away from an active wild-dog denning site. The top-panel shows wild dog utilization during the pre-denning period (Sep–Nov 2011) while the bottom panel shows the same area during the active denning period (Dec 2011–Jan 2012). White lines are control road transects and black lines are the 'near den' (i.e., inside the denning home range) transects. We used a length weighted mean utilization of 0.005 to partition the denning home range and control areas. Points show GPS relocations (30 min interval) of a single adult female from this pack during the crepuscular hunting period. There are no GPS relocations available in the post-denning period but tracking VHF collars fitted to other members of the same pack confirmed that the den was not being used during this time.

5 km



FIG. B2. Monthly rainfall during each year of this study. Wild dogs recolonized the area in 2002 and began denning on Mpala Research Centre in 2003. Shaded sections show the years during which the two herbivore exclosure experiments were conducted. We estimated dik-dik abundance in 1999–2002 and from 2008-onwards.



FIG. B3. The effect of wild dog recovery and ungulate exclusion on tree abundance, measured as the net rate of change in density (individuals 100 m<sup>-2</sup> yr<sup>-1</sup>) for trees in the 1–2 m height class, over a 3-year period pre- (1999–2002, n = 3 pairs of plots) and post- (2009–2012, n = 6 pairs of plots) wild dog recovery. Treatments were either TOTAL (no access to ungulates) or OPEN (accessible to all ungulates). "Community" includes the aggregate response of 32 species of trees and woody plants.

TABLE B1. Model selection table for the effects of dik-dik abundance (DENSITY = individuals km<sup>-2</sup>), rainfall (RAIN = mm·6 mo<sup>-1</sup>), the estimated consumption of dik-dik by wild dogs prior to dik-dik population surveys (EATEN = individuals·km<sup>-2</sup>) and an interaction between EATEN and DENSITY and RAIN and DENSITY on dik-dik population growth (*rd*) between surveys. *K* refers to the number of parameters in the model.

Model	K	Log- likelihood	AICc	ΔAICc	Akaike weight (wi)
EATEN	6	-66.66	155.83	0.00	0.92

EATEN + RAIN	7	-66.06	162.12	6.29	0.04
EATEN + DENSITY	7	-66.66	163.33	7.50	0.02
RAIN	6	-71.29	165.08	9.25	0.01
DENSITY	6	-71.45	165.41	9.58	0.01
EATEN + RAIN + EATEN $\times$ RAIN	8	-65.62	171.24	15.41	<0.01
DENSITY + RAIN	7	-74.12	178.23	22.40	<0.01
EATEN + DENSITY + EATEN $\times$ DENSITY	8	-73.30	186.60	30.77	<0.01
EATEN + DENSITY + RAIN	8	-74.35	188.70	32.87	<0.01
EATEN + DENSITY + RAIN + EATEN × DENSITY	9	-69.59	193.17	37.34	<0.01
EATEN + DENSITY + RAIN + EATEN × RAIN	9	-71.42	196.83	41.00	<0.01
EATEN + DENSITY + RAIN + EATEN × DENSITY + EATEN × RAIN	10	-62.17	199.34	43.51	<0.01

TABLE B2. Model selection table for the effects of ungulate exclusion treatment (OPEN vs. TOTAL exclusion) and the status of wild dog recovery (pre- vs. post-wild dog recovery) on tree abundance, measured as the net difference in stem density (individuals  $\cdot 100 \text{ m}^{-2} \cdot \text{yr}^{-1}$ ). "Community" includes the aggregate response of 32 species of trees and woody plants. *K* refers to the number of parameters in the model.

Species	Model	K	Log- likelihood	AICe	ΔΑΙCc	Akaike weight (wi)
Community	Treatment + Recovery	7	-3.65	32.51	0.00	0.75
	Recovery	6	-7.97	35.57	3.06	0.16
	Treatment	6	-9.00	37.63	5.12	0.06
	Treatment + Recovery + Treatment × Recovery	8	-3.65	39.30	6.80	0.03
A. etbaica	Treatment	6	26.99	-34.33	0.00	0.56

	Treatment + Recovery	7	29.47	-33.74	0.59	0.41
	Treatment + Recovery + Treatment × Recovery	8	30.00	-28.00	6.33	0.02
	Recovery	6	22.62	-25.61	8.72	0.01
A. mellifera	Treatment + Recovery	7	24.07	-22.94	0.00	0.37
	Treatment	6	21.29	-22.94	0.00	0.37
	Recovery	6	20.89	-22.14	0.80	0.25
	Treatment + Recovery + Treatment × Recovery	8	24.23	-16.46	6.48	0.01
<i>Grewia</i> spp.	Treatment	6	23.96	-28.29	0.00	0.67
	Treatment + Recovery	7	25.95	-26.71	1.58	0.30
	Treatment + Recovery + Treatment × Recovery	8	26.94	-21.88	6.40	0.03
	Recovery	6	18.75	-17.86	10.43	< 0.01

TABLE B3. Changes in the energetic demand of browsers at Mpala Research Centre pre- (1999–2002) and post- (2008–2011) recovery of wild dogs. Population densities were determined using distance sampling methods, described in <u>Appendix A</u>.

			Density (individuals • km <sup>-2</sup> )		Biomass density (kg · km <sup>-2</sup> )			<sup>7</sup> Energetic demand (KJ · km <sup>-2</sup> )			
Species	Mass (kg)	Residency	Pre-	Post	-	Pre-	Post-		Pre-		Post-
Dik-dik	5	Resident	139.00	109.00	694		546	9340	6	78	359
Steinbuck	10	Resident	0.52	0.10*	5		1	2574		76	7
Thompson's gazelle	25	Resident	0.00	0.04*	0		4 0			21	23
Bushbuck	30	Resident	0.20	0.10*	0.10* 6		3	2859		17	19

Impala	40	Resident	20.30	23.30	812	932	104872	116037
Grant's gazelle	50	Resident	0.00	1.35	0	68	0	16895
Eland	340	Resident	0.37	0.39	126	133	26682	27733
Giraffe	750	Migrant	0.33	1.56	248	1170	43846	137118
Elephant	1725	Migrant	1.70	0.98	2933	1691	269153	179643
All browsers			162	137	4823	4547	543392	560395
Resident-only				135	1643	1686	230393	243634

\* Due to low detection, these estimates are at the limit of resolution.

[Back to E096-238]